

ONCOLOGY KNOWLEDGE HUB · FOR HEALTH PROFESSIONALS

Colon Cancer

A mini review for clinicians (from the Colorectal Cancer module)

For postgraduate residents (medical, surgical and radiation oncology, general surgery, gastroenterology, medicine), practising clinicians and medical students

Dr. Haritha Chiramana

MD (Internal Medicine), DM (Medical Oncology)

Senior Medical Oncologist & Hemato-Oncologist

C.R. Reddy Cancer Center, C.R. Reddy Superspeciality Hospital

17/406, Old Post Office Road, near Leela Mahal Road, Nellore, Andhra Pradesh 524003

Phone 0861 232 5344 · WhatsApp +91 79896 42272

info@drharitha.co.in · drharitha.co.in

Dr. Haritha teaches oncology to postgraduates and medical students. Colleges and departments may write to info@drharitha.co.in for teaching sessions.

CONTENTS

- | | |
|---|--|
| 01 At a glance | 07 Management algorithms |
| 02 Clinical presentation, common traps & genetic causes | 08 Survivorship, follow-up & reducing recurrence |
| 03 Screening, workup & staging essentials | 09 Landmark trials |
| 04 Biomarkers that change treatment | 10 Practice pearls & pitfalls |
| 05 Local therapy: surgery & radiotherapy | 11 Case vignettes: lines of management |
| 06 Systemic therapy: indications & options | 12 References & guidelines |

Cite as: Chiramana H. Colon Cancer: a mini review for clinicians. Oncology Knowledge Hub [Internet]. Nellore (India): drharitha.co.in; 2026 [updated 2026 Oct 9]. Available from: <https://drharitha.co.in/professionals/colorectal-cancer> · Free for education with attribution (CC BY-NC-ND 4.0).

Updated 9 October 2026. Educational summary written in our own words from current guidelines and primary trials; no drug doses; not advice for an individual patient. Verify current guidelines and Indian regulatory approvals before treatment decisions. Patients and families: please see the patient guides at drharitha.co.in.

Colon Cancer

From the Colorectal Cancer module · colon and metastatic disease · Key points for clinicians · Last updated 9 October 2026

01 At a glance

- Colorectal cancer is the third commonest cancer in the world and the second commonest cause of cancer death. India records roughly 65,000 new cases a year (GLOBOCAN 2022 estimates); age-standardised rates are lower than in the West but rising in cities, and Indian hospital series report a larger share of **rectal cancers** and of patients **under 50**, with more signet-ring and mucinous histology.
- Most cancers arise from a visible polyp over about a decade, so **colonoscopy prevents cancer as well as finding it**. India has no organised colorectal screening programme; a stool immunochemical test (FIT) is the practical first step for average-risk adults.
- Stage drives survival: about 90% of patients with localised disease are alive at 5 years, about 70% with nodal disease, and about 15% with distant metastases. Even so, selected patients with liver or lung metastases are **cured by resection**.
- Four decisions shape treatment: **site** (colon or rectum, and right or left), **stage** (CT for everyone, MRI pelvis for the rectum), **mismatch-repair status** (every new cancer) and, when metastatic, **RAS, BRAF and HER2**.
- The commonest avoidable harm in Indian practice is delay: rectal bleeding labelled as piles, anaemia treated with iron alone, and caecal masses treated as intestinal TB without tissue.

Approximate share by site; Indian hospital series often report a larger rectal share and more patients under 50 than Western series.

Right colon · ~30%	Left colon & sigmoid · ~30%	Rectum · ~40%
Right colon <i>Caecum to transverse colon</i> - Anaemia, fatigue, a mass; bleeding is occult - Older, more often women; more MSI-high and BRAF - Poorer prognosis when metastatic Metastatic, RAS/BRAF wild-type: doublet + bevacizumab	Left colon & sigmoid <i>Splenic flexure to sigmoid</i> - Change in bowel habit, bleeding, obstruction - Chromosomal-instability type; HER2 more often - Better prognosis when metastatic Metastatic, RAS/BRAF wild-type: doublet + anti-EGFR	Rectum <i>Up to 15 cm from the anal verge</i> - Bleeding, tenesmus, mucus; often called "piles" - Staged locally by MRI pelvis - Pelvic recurrence risk: radiotherapy matters Total neoadjuvant therapy; organ preservation for some

Figure 1. Three working sites. Where the tumour starts changes the symptoms, the staging (MRI for the rectum), the role of radiotherapy and, when metastatic, the choice of biological drug.

02 Clinical presentation, common traps & genetic causes

Who is at risk: ask about each of these

Factor	Approximate size of risk	Clinical note
Age	Risk rises steeply after 45–50	Early-onset cancer (under 50) is rising worldwide; Indian series often report a fifth to a third under 45. Symptoms in the young must not be dismissed.
One first-degree relative with colorectal cancer	About 2-fold; higher if diagnosed before 50 or several relatives	Start colonoscopy at 40, or 10 years before the youngest relative's diagnosis.
Lynch syndrome	Lifetime risk roughly 10–50% depending on the gene (highest MLH1/MSH2)	Also endometrial, ovarian, gastric, urothelial and small-bowel cancers. Section 04.
Polypsis syndromes (FAP, MUTYH, serrated polyposis)	Classic FAP near 100% without colectomy	Hundreds of adenomas from the teens; attenuated forms present later.
Inflammatory bowel disease	Rises with duration and extent of colitis	Surveillance colonoscopy with chromoendoscopy from about 8 years after onset of extensive colitis.
Primary sclerosing cholangitis (PSC) with IBD	About 3-fold higher than IBD alone (meta-analysis^[1]); cancers are often right-sided, at a median age near 50	Colonoscopy at PSC diagnosis (colitis is often silent: take random biopsies), then every year ; PSC without IBD: every 5 years. Ursodeoxycholic acid is not a chemopreventive.
Lifestyle	Each modest (relative risks 1.1–1.5)	Red and processed meat, obesity, inactivity, smoking, alcohol and diabetes raise risk; fibre, calcium and physical activity lower it.
Previous adenoma or colorectal cancer	Metachronous cancer about 1–2% over 5 years	Surveillance colonoscopy (section 08).
Pelvic radiotherapy, acromegaly, cystic fibrosis, Streptococcus gallolyticus bacteraemia	Increased	Colonoscopy after S. gallolyticus endocarditis or bacteraemia.

How it presents

Site	Typical symptoms	Clue that is often missed
Right colon	Fatigue, iron-deficiency anaemia, right iliac fossa mass, vague pain	Bleeding is occult: anaemia in a man or post-menopausal woman needs both endoscopies.
Left colon and sigmoid	Change in bowel habit, bright or dark blood mixed with stool, colicky pain, obstruction	Alternating constipation and loose stools; “pencil” stools.
Rectum	Bleeding, tenesmus, mucus, incomplete evacuation, pain at the anus if low	A digital rectal examination finds most low tumours; it is skipped surprisingly often.
Emergency (about 1 in 5)	Large-bowel obstruction, perforation, massive bleeding	Emergency presentation predicts worse survival; urgent CT, not repeated enemas.




Figure 2. Rectal bleeding and anaemia: piles or cancer? The examination comes first, but persistent symptoms, anaemia or a positive FIT need colonoscopy whatever the age.

- **Urgent referral triggers** (2-week pathway): rectal bleeding with a change in bowel habit; unexplained iron-deficiency anaemia; a rectal or abdominal mass; a positive FIT in a symptomatic person; persistent change in bowel habit over 40–45; weight loss with abdominal symptoms. Same day: obstruction, perforation or heavy bleeding.
- **FIT in symptomatic patients:** a faecal haemoglobin at or above 10 µg/g identifies most cancers and is used in the UK to prioritise colonoscopy; a negative FIT does not override a mass, anaemia or rectal bleeding.
- **Atypical presentations:** fever of unknown origin, a malignant fistula, *Streptococcus gallolyticus* (bovis) bacteraemia or endocarditis, *Clostridium septicum* sepsis, or adenocarcinoma of unknown primary – each warrants colonoscopy.
- **CEA is not a diagnostic test:** it is normal in many early cancers and raised in smokers, liver disease and inflammation. Use it for baseline and follow-up only.
- **Ileocaecal TB versus cancer:** both cause a right iliac fossa mass, anaemia and weight loss. Take multiple biopsies for histology, TB PCR and culture; empirical antituberculous treatment without proof can hide a caecal cancer for months.

Hereditary syndromes to recognise

Syndrome	Gene	Clues	What to do
Lynch syndrome	MLH1, MSH2 (EPCAM), MSH6, PMS2	dMMR tumour; young; right colon; endometrial, ovarian, gastric, urothelial and small-bowel cancers in the family	Colonoscopy every 1–2 years from 20–25 (MLH1/MSH2) or every 1–3 years from 30–35 (MSH6/PMS2), or 2–5 years before the youngest family case; upper endoscopy with H. pylori treatment from 30–35; daily aspirin recommended for MLH1, MSH2, MSH6 and EPCAM carriers (CAPP2^[2]), individualised for PMS2
Familial adenomatous polyposis	APC	>100 adenomas, often in the teens; desmoids, duodenal adenomas	Prophylactic colectomy; upper GI surveillance

Syndrome	Gene	Clues	What to do
MUTYH-associated polyposis	MUTYH, biallelic (recessive)	Usually up to 100 adenomas by the 50s–60s, but sometimes cancer with few or no polyps; serrated polyps too; siblings affected, parents not	Colonoscopy every 1–2 years from 25–30; upper endoscopy from 30–35 (interval by Spigelman stage); yearly thyroid examination and ultrasound. Single-allele carriers (1–2% of people) screen as the general population (Win 2014^[3]); siblings have a 25% chance of being affected
Polymerase proofreading-associated polyposis (PPAP)	POLE, POLD1 exonuclease domain (dominant)	Typically 30–100 adenomas; colorectal cancer risk over 20%; duodenal adenomas; possibly endometrial cancer	Colonoscopy from 25–30 every 2–3 years (1–2 years once polyps appear); upper endoscopy from 25–30 (POLE) or 30–35 (POLD1) about every 3 years. Tumours are hypermutated and respond to immune checkpoint inhibitors (Palles 2022^[4])
Serrated polyposis	Usually none found	Many serrated polyps, right colon	Clearance colonoscopy; relatives screened earlier
Peutz–Jeghers, juvenile polyposis	STK11; SMAD4/BMPRIA	Pigmented lips; hamartomas; HHT with SMAD4	Specialist surveillance of several organs

- Refer for genetic counselling when there is a dMMR tumour not explained by MLH1 methylation or BRAF, 10 or more adenomas, cancer under 50, or a strong family history. About one in six patients diagnosed under 50 carries a germline predisposition, so a germline panel for everyone diagnosed under 50 is reasonable where affordable.
- Family history alone misses most Lynch carriers, although about one in four patients with colorectal cancer has an affected relative. A variant of uncertain significance (VUS) does not change management: manage by the personal and family history and ask the laboratory for re-classification about once a year.

SHARE WITH PATIENTS: SYMPTOMS AND EARLY DIAGNOSIS

Plain-language myth pages written for patients and families. Send them from the clinic to reinforce what you discussed.

[Telugu–English handout: bowel cancer warning signs](#)
(తెలుగు · English)

[Blood in the stools is always caused by piles?](#)

[Colorectal cancer occurs only in elderly people?](#)

[If I am anaemic, iron tablets are all I need?](#)

[A lump in the right side of the abdomen is usually intestinal TB?](#)

[If I have no digestive problems, I can't have colorectal cancer?](#)

03 Screening, workup & staging essentials




Figure 3. How colorectal cancer develops. Most cancers pass through a visible polyp stage, which is why colonoscopy can prevent cancer as well as find it. The route also predicts MMR status and BRAF, both of which change treatment.

Screening options for average-risk adults

Test	Evidence	Practical points
FIT (faecal immunochemical test), yearly (one sample)	Guaiac stool testing cut colorectal cancer deaths by about a third with annual testing (Minnesota FOBT trial^[5]); FIT is more sensitive and needs no diet change	Best suited to Indian practice: cheap, one sample, no preparation. One-time sensitivity for cancer is about 74% (lower for advanced adenomas), so yearly repetition matters. Return the sample within about 24 hours and keep it away from heat — delay and high temperatures reduce sensitivity, a real issue in Andhra summers. A positive result needs colonoscopy within weeks; it is not a test to repeat.
Colonoscopy every 10 years	Removing adenomas roughly halved colorectal cancer deaths (National Polyp Study^[6]). In NordICC^[7] an invitation to one colonoscopy reduced incidence by about a fifth at 10 years; the death reduction was not significant because only 42% attended	Diagnostic and therapeutic; quality matters (adenoma detection rate, caecal intubation, good bowel preparation). Small risk of bleeding and perforation.
CT colonography every 5 years	Sensitive for polyps ≥10 mm	Option when colonoscopy is incomplete or refused; findings still need colonoscopy.
Flexible sigmoidoscopy	Reduced distal cancer incidence and mortality in randomised trials	Misses right-sided lesions; less used now.
Multitarget stool DNA (with FIT), every 1–3 years	More sensitive than FIT for cancer; benefit on outcomes is indirect	A reasonable alternative where available; costly in India.
Blood-based cfDNA tests, stool RNA tests	Efficacy data not yet robust; poor sensitivity for advanced adenomas	Not recommended for screening at present. Never use a single in-office guaiac test after a rectal examination, or a barium enema.


- **When to start and stop:** screen average-risk adults from 45 to 75 ([USPSTF 2021^{\[8\]}](#) lowered the start age to 45; benefit is clearest from 50); individualise from 76 to 85 by health and previous results (screening up to

85 is reasonable if never screened). Colonoscopy every 10 years or yearly FIT are the preferred options; whatever the test, adherence matters more than the choice, and every positive non-colonoscopy test needs a prompt colonoscopy.

- **Family history:** one first-degree relative with colorectal cancer, an advanced adenoma or an advanced serrated lesion → colonoscopy from 40 (or 10 years before the relative's diagnosis, whichever is earlier), repeated every 5 years; yearly FIT if colonoscopy is declined. Ask every adult about family history before 40 and update it every few years.
- **An Indian approach:** opportunistic FIT for adults over 45–50 seen in clinics, colonoscopy for a positive FIT, symptoms, anaemia or a family history; quality colonoscopy capacity is the main constraint.
- **Polyps:** adenomas (risk rises with size, villous histology and high-grade dysplasia), sessile serrated lesions and traditional serrated adenomas are precancerous; small distal hyperplastic polyps are not. Inflammatory polyps need no removal unless symptomatic. Juvenile polyposis and Peutz–Jeghers syndrome carry cancer risk.
- **Malignant polyp** (cancer invading the submucosa in a removed polyp): no further surgery if it was removed in one piece with a clear margin (≥ 1 mm), grade 1–2, no lymphovascular invasion and low tumour budding; otherwise resection with nodes. Tattoo the site.

Workup before treatment: the checklist

Step	Why	Notes
Complete colonoscopy with biopsies and tattoo	Histology, synchronous cancers (3–5%) and polyps	If obstructing: CT (or MR) colonography — preferred over barium enema — or complete colonoscopy within 3–6 months after surgery.
CT chest, abdomen and pelvis with contrast	Distant staging; local extent of colon cancer	Radiological T4 or bulky T3 colon cancer may suit neoadjuvant chemotherapy (section O6).
MRI liver (with hepatocyte-specific contrast if available)	Indeterminate or potentially resectable liver lesions	Before any liver surgery or ablation.
MRI pelvis, rectal protocol (rectal cancer)	Local stage: T, CRM, EMVI, nodes, height, sphincter	Done before any treatment; repeat after neoadjuvant therapy. Endorectal ultrasound adds detail for early T1–T2 tumours.
CEA, before and after surgery	Baseline for follow-up; a high level carries a worse prognosis	Not for screening or diagnosis.
PET-CT	Selected cases only	Before resecting metastases, or for a rising CEA with normal CT. Not routine for staging.
Blood counts, liver and kidney function; DPYD genotype where available	Fitness and safety of chemotherapy	Correct anaemia; nutritional assessment; geriatric assessment over 70.
MMR immunohistochemistry on the biopsy	Changes treatment at every stage; screens for Lynch	Request at diagnosis so neoadjuvant options are not missed.

Depth of invasion (T)**N (regional nodes; examine ≥12)**

N1a: 1 node · N1b: 2–3 nodes
 N1c: tumour deposits, nodes negative
 N2a: 4–6 nodes · N2b: 7 or more

M (distant)

M1a: one organ
 M1b: two or more organs
 M1c: peritoneum (± organs)

Stage groups


I: T1–2 N0 · IIA T3 · IIB T4a · IIC T4b
 III: any T, N1–2 (IIIA–C)
 IV: any M1 (IVA–C)

Figure 4. TNM at a glance (AJCC/UICC 8th edition). A pathology report with fewer than 12 lymph nodes may understage the cancer; tumour deposits without nodes count as N1c.

- **Obstructing left-sided cancer:** emergency resection (Hartmann's or primary anastomosis with or without a stoma) or a self-expanding stent as a bridge to elective surgery a few weeks later in expert hands; stents are avoided if bevacizumab is planned.
- **Prognosis within stage III** varies widely: T1–2 N1 does as well as many stage II tumours, while T4 N2 does worse than some stage IV patients after metastasectomy.

04 Biomarkers that change treatment

- **Mismatch repair (MMR) / microsatellite instability (MSI):** test every new cancer by immunohistochemistry (MLH1, MSH2, MSH6, PMS2) or PCR/NGS. About 15% of all colorectal cancers are dMMR (more in stage II, right colon), but only about 4% of metastatic ones. dMMR changes adjuvant treatment (no fluoropyrimidine alone in stage II), neoadjuvant treatment (immunotherapy is highly active) and metastatic treatment (immunotherapy first line).
- **RAS (KRAS and NRAS, exons 2, 3 and 4):** mutated in about 45–50%. Any RAS mutation excludes anti-EGFR antibodies, which can harm these patients ([PRIME^{\[9\]}](#)). Test all of KRAS and NRAS, not just KRAS exon 2.
- **KRAS G12C** (about 3%): sotorasib or adagrasib combined with an anti-EGFR antibody after chemotherapy ([CodeBreak 300^{\[10\]}](#)); alone they work poorly in colorectal cancer because EGFR signalling bypasses the block.
- **BRAF V600E** (about 8–10%): poor prognosis, right-sided, peritoneal spread. Encorafenib with cetuximab and FOLFOX in first line ([BREAKWATER^{\[11\]}](#)), encorafenib with cetuximab later ([BEACON^{\[12\]}](#)). Non-V600 BRAF mutations behave differently.
- **HER2 amplification** (about 3–5%, mostly RAS wild-type, left-sided): tucatinib with trastuzumab ([MOUNTAINEER^{\[13\]}](#)) or trastuzumab deruxtecan ([DESTINY-CRC01^{\[14\]}](#)). Use colorectal-specific scoring; HER2 also predicts resistance to anti-EGFR antibodies.
- **POLE / POLD1 proofreading mutations** (somatic or germline): ultramutated tumours (often >50 mutations per megabase) that respond to immune checkpoint inhibitors as dMMR tumours do; immunotherapy is suggested first line for such metastatic cancers.
- **PI3K-pathway alterations** (*PIK3CA* exon 9/20 and other *PIK3CA*, *PIK3R1* or *PTEN* mutations; roughly a third of cancers): after resection of stage I–III disease, low-dose aspirin for 3 years roughly halved recurrence in [ALASCCA^{\[15\]}](#). This makes NGS of the resected tumour useful even in early disease; aspirin can start once the patient has recovered from surgery and run alongside adjuvant chemotherapy.
- **Rare fusions** (NTRK, RET, ALK; under 1%, mainly MSI-high, RAS/BRAF wild-type): tumour-agnostic TRK or RET inhibitors.
- **Primary tumour side** works like a biomarker in metastatic RAS/BRAF wild-type disease: left-sided tumours gain survival from anti-EGFR antibodies ([PARADIGM^{\[16\]}](#)), right-sided ones generally do not ([pooled analysis^{\[17\]}](#)).
- **Liquid biopsy (ctDNA):** useful when tissue is scant (RAS concordance with tissue is high in most patients), for anti-EGFR rechallenge (RAS wild-type in plasma again), and after surgery to estimate relapse risk ([DYNAMIC^{\[18\]}](#)) — the latter still largely within trials and costly.
- **Germline pharmacogenomics:** DPYD variants (severe fluoropyrimidine toxicity) and UGT1A1*28 (irinotecan) — test before treatment where available.
- **Not useful for treatment choice:** CEA level, consensus molecular subtypes, immunoscore (prognostic only), and tumour mutational burden outside MSI-high.



Family history alone misses about half of Lynch carriers. Somatic “double hits” (two tumour mutations) explain many unexplained dMMR cases, but germline testing still comes first.

Figure 5. One test, three uses. MMR status changes adjuvant, neoadjuvant and metastatic treatment, and it is the screening test for Lynch syndrome.

Approximate frequency in metastatic colorectal cancer (overlaps exist; NTRK, RET and others are each under 1%).





Figure 6. Biomarkers that change treatment. Each result points to a specific drug, or away from one: anti-EGFR antibodies help only RAS and BRAF wild-type tumours, ideally left-sided.

05 Local therapy: surgery & radiotherapy

Colon cancer surgery


- **Principle:** remove the bowel segment with its mesentery and the lymph nodes along the feeding vessels, with clear margins (usually 5 cm or more of bowel on each side) and at least 12 nodes for staging.
- **Complete mesocolic excision with central vascular ligation** (CME/CVL, “D3”) removes the mesocolon in its fascial envelope and the central nodes; it yields more nodes and may lower recurrence in observational series, at the cost of more vascular injury if done without experience.
- **Laparoscopic versus open:** equivalent long-term cancer outcomes in [COST^{\[19\]}](#) and [COLOR^{\[20\]}](#), with faster recovery; robotic surgery is an option where available. Enhanced recovery (ERAS) pathways shorten stay and complications.
- **T4b tumours** (adherent to another organ): resect en bloc; separating the adhesion risks tumour spill. Consider neoadjuvant chemotherapy first.
- **Synchronous cancers or polyposis:** subtotal or total colectomy; Lynch patients under about 60 may choose extended colectomy to reduce metachronous risk.



Operation	Best suited to	Advantages	Disadvantages
Right / extended right hemicolectomy	Caecum, ascending, hepatic flexure, transverse colon	Safe ileocolic anastomosis; good node yield	Looser stools for some; middle colic dissection in extended resections

Operation	Best suited to	Advantages	Disadvantages
Left hemicolectomy / sigmoid colectomy	Descending and sigmoid colon	Bowel function usually normal afterwards	Splenic flexure mobilisation; anastomotic leak 2–5%
Subtotal / total colectomy	Synchronous cancers, Lynch (selected), obstructing left cancer with caecal damage	Removes the at-risk colon	Frequent stools
Low anterior resection with TME	Mid and low rectum when the sphincter can be preserved	Avoids a permanent stoma	Low anterior resection syndrome; temporary ileostomy; leak 5–15%
Abdominoperineal resection	Tumours invading the sphincter or levators; poor continence	Clear distal margin	Permanent colostomy; perineal wound problems
Local excision (TEM / TAMIS)	Selected cT1 low-risk rectal tumours; after a good response in trials	No stoma; minimal function loss	Nodes not removed: higher local recurrence unless the tumour is very favourable
Hartmann's operation	Emergency obstruction or perforation; frail patients	No anastomosis at risk	Stoma; reversal often never done

Local therapy for metastases



- **Liver resection** is the only curative option for liver metastases: 5-year survival after resection now approaches 50–60% in modern series, with operative mortality under 5% (under 1% in high-volume centres). Resectability depends on what remains — inflow, outflow, biliary drainage and a future liver remnant above 20% (above 30% after chemotherapy-related liver injury) — not on the number of lesions. Aim for margins over 10 mm, but resect if more than 1 mm is achievable; parenchyma-sparing resections are favoured, with wider margins for RAS- or BRAF-mutant tumours. Repeat hepatectomy is feasible for recurrence.
- **Sequencing with chemotherapy:** fit patients with up to 4 lesions in one lobe and RAS/BRAF wild-type tumours usually go straight to surgery; higher-risk disease (more than 4 lesions, bilobar, RAS/BRAF-mutant,


suspicious portal nodes or extrahepatic disease) has chemotherapy first. Perioperative FOLFOX improved progression-free but not overall survival in [EPOC](#)^[21] ([long-term results](#)^[22]). Give no more than about 4 cycles before surgery, re-image every 6–8 weeks, operate as soon as lesions become resectable (at least 4 weeks after chemotherapy, 6–8 weeks after bevacizumab), and complete about 6 months of treatment in total. Mark small (<2 cm) or deep (>1 cm from the surface) lesions with fiducials before chemotherapy: they may vanish on imaging but rarely disappear pathologically, and should still be resected.

- **Conversion therapy** for initially unresectable liver-only disease: RAS/BRAF wild-type → oxaliplatin-based doublet or triplet (with an anti-EGFR antibody for left-sided tumours); RAS- or BRAF-mutant and fit → FOLFOXIRI with bevacizumab, which raised resection rates in [CAIRO5](#)^[23]. Even dMMR potentially resectable disease is usually treated with chemotherapy rather than upfront immunotherapy, although this is evolving. Hepatic arterial infusion remains a trial or expert-centre option.
- **Synchronous metastases**: a bleeding, obstructing or perforated primary is resected first; extensive bilobar disease suits a planned two-stage hepatectomy with portal vein embolisation; a right-colon primary with limited liver disease can be removed in one operation (liver first); a reverse (liver-first) approach is an option when the liver disease dominates.
- **Ablation and other liver-directed therapy**: for up to a few lesions of 3 cm or less, thermal ablation (radiofrequency or microwave) gave similar survival with fewer complications than resection in [COLLISION](#)^[24] (long-term follow-up awaited); SBRT, irreversible electroporation, TACE or yttrium-90 radioembolisation suit selected unresectable or chemotherapy-refractory liver-predominant disease. Adding radioembolisation to first-line chemotherapy did not improve survival ([FOXFIRE/SIRFLOX combined analysis](#)^[25]).
- **Lung**: resection or SBRT for limited, slow-growing lung metastases; long-term survival is common when the disease-free interval is long.
- **Other sites**: selected isolated retroperitoneal nodal, adrenal or bulky symptomatic ovarian metastases can be resected; multi-organ disease responding to chemotherapy is not debulked (no survival gain, more toxicity).
- **Peritoneum**: complete cytoreductive surgery in expert centres for limited disease without extraperitoneal spread (low peritoneal cancer index), after shared decision-making about its morbidity; adding oxaliplatin HIPEC gave no survival benefit and more late complications in [PRODIGE 7](#)^[26].
- **The primary tumour in stage IV**: in patients with unresectable metastases and an asymptomatic primary, upfront resection did not improve survival ([JCOG1007](#)^[27]; [SYNCHRONOUS/CCRe-IV](#)^[28]); resect, bypass, divert or stent only for obstruction, bleeding or perforation. With an asymptomatic primary and resectable metastases, systemic therapy usually comes first so that the biology declares itself.
- **Liver transplant**: for highly selected unresectable liver-only disease after good response to chemotherapy, transplant plus chemotherapy improved 5-year survival to about 57% versus 13% with chemotherapy alone ([TRANSMET](#)^[29]); needs organ allocation policy and expert centres.

o6 Systemic therapy: indications & options

Indications and options only; no doses. Benefits are given in absolute terms where possible. Trial names link to the primary publications.

Adjuvant therapy after resection of colon cancer




- **Stage III:** adding oxaliplatin to a fluoropyrimidine raised 6-year overall survival by about 4 percentage points ([MOSAIC](#)^[30], [OS update](#)^[31], [NSABP C-07](#)^[32]); capecitabine can replace infusional 5-FU ([X-ACT](#)^[33]). Start within about 8 weeks of surgery; late starts lose benefit.
- **Duration** ([IDEA](#)^[34]): three months of CAPOX is enough for low-risk stage III (T1–3 N1), with much less lasting neuropathy; high-risk stage III (T4 or N2) does better with 6 months, especially with FOLFOX. If oxaliplatin causes troublesome neuropathy, stop it and continue the fluoropyrimidine.
- **Stage II:** the absolute gain from chemotherapy is about 3–4% ([QUASAR](#)^[35]). Offer it for high-risk features (T4, fewer than 12 nodes, obstruction or perforation, lymphovascular or perineural invasion, poorly differentiated pMMR tumours, close margins, high tumour budding). Low-risk pMMR: discuss, often observe. dMMR stage II: no fluoropyrimidine alone (no benefit; may harm); T4 dMMR can be considered for oxaliplatin-based treatment.
- **Stage III dMMR:** adding atezolizumab to 6 months of FOLFOX, continued alone to 12 months, roughly halved the risk of recurrence or death ([ATOMIC](#)^[36]); overall survival did not differ at 5 years, and grade 3–4 toxicity was higher. Where atezolizumab is not accessible, standard oxaliplatin-based adjuvant therapy remains appropriate.
- **Older patients:** oxaliplatin adds little over the age of 70; a fluoropyrimidine alone is reasonable for many. Use geriatric assessment rather than age alone.
- **Adjunctive aspirin:** in unselected patients, randomised trials of aspirin ([ASCOLT](#)^[37]) and celecoxib ([CALGB/SWOG 80702](#)^[38]) after adjuvant therapy did not improve disease-free or overall survival, despite favourable observational data; discuss it, or reserve it for those with another indication. For stage I–III

cancers with a PI3K-pathway alteration, low-dose aspirin for 3 years reduced recurrence (about 8% vs 14–17%) in [ALASCCA](#) [15] (celecoxib also helped the *PIK3CA*-mutant subgroup of [80702](#) [39], but aspirin is safer).

- **Vitamin D:** low levels are linked with worse outcomes, but supplementation has not been proven to improve survival ([meta-analysis of trials](#) [40]). It is reasonable to measure 25-hydroxyvitamin D at diagnosis and replace it when below 20 ng/mL, mainly for bone health.
- **ctDNA-guided treatment:** in stage II, giving chemotherapy only to ctDNA-positive patients halved its use without losing recurrence-free survival ([DYNAMIC](#) [18]). Promising, but not yet standard and expensive in India.
- **Neoadjuvant chemotherapy for colon cancer:** for radiologically locally advanced (T4 or bulky T3) pMMR tumours, 6 weeks of FOLFOX before surgery reduced 2-year residual or recurrent disease from about 22% to 17% and downstaged tumours ([FOX TROT](#) [41]). dMMR tumours respond poorly to this chemotherapy but extraordinarily to immunotherapy: two cycles of nivolumab with ipilimumab gave a major pathological response in 95% and a complete response in two-thirds ([NICHE-2](#) [42]). For dMMR tumours that are locally unresectable, immunotherapy is an appropriate alternative to chemotherapy, ideally in a trial; most resectable colon cancers still go straight to surgery.

Metastatic disease



- **First line, pMMR, RAS/BRAF wild-type, left-sided:** FOLFOX or FOLFIRI with panitumumab or cetuximab improved overall survival over bevacizumab in left-sided tumours ([PARADIGM](#) [16], [FIRE-3](#) [43], [CALGB/SWOG 80405](#) [44], [CRYSTAL](#) [45]). Response rates are highest, which matters when conversion to resection is the aim.
- **Right-sided or RAS-mutant:** doublet chemotherapy with bevacizumab ([Hurwitz](#) [46]); for a right-sided RAS/BRAF wild-type tumour where bevacizumab is contraindicated, use chemotherapy alone rather than an anti-EGFR antibody. For fit patients with a large tumour burden, FOLFOXIRI with bevacizumab raises response and survival at the price of more toxicity ([TRIBE](#) [47]); limit the triplet to 3–6 months, then fluoropyrimidine maintenance.

- **Choosing the backbone:** FOLFOX, CAPOX and FOLFIRI are equally effective first line; choose by toxicity, and prefer FOLFIRI if adjuvant oxaliplatin was given within the past 12 months. Start chemotherapy early, before symptoms, rather than waiting. Stop oxaliplatin after 3–4 months (before disabling neuropathy) and continue the rest; planned breaks are reasonable for responding, low-volume disease. Judge response by imaging: a rising CEA alone should be confirmed radiologically before changing treatment.
- **dMMR / MSI-high (and POLE/POLD1-ultramutated):** immunotherapy first. Nivolumab with ipilimumab was superior to chemotherapy and to nivolumab alone ([CheckMate 8HW](#)^[48]; [second report](#)^[49]) and is preferred for fit patients; pembrolizumab ([KEYNOTE-177](#)^[50], which doubled progression-free survival over chemotherapy), nivolumab or dostarlimab alone suit less fit patients. Continue for 1–2 years if responding; stop after at least a year if a complete response is confirmed on two scans. On progression, switch to fluoropyrimidine-based chemotherapy ± a biological agent.
- **BRAF V600E:** encorafenib + cetuximab + mFOLFOX6 roughly doubled median overall survival over standard first-line chemotherapy (about 30 vs 15 months; [BREAKWATER](#)^[11]), whatever the side; FOLFOX is the preferred backbone, FOLFIRI an alternative with less neuropathy. After first line without it, encorafenib + cetuximab improved survival over chemotherapy ([BEACON](#)^[12]); after progression on the triplet, FOLFIRI or irinotecan with bevacizumab. Anti-EGFR antibodies alone do not work in BRAF-mutant disease.
- **Maintenance:** after 4–6 months of induction, de-escalate to a fluoropyrimidine with bevacizumab ([CAIRO3](#)^[51]) or with anti-EGFR; full treatment breaks suit selected patients. Reintroduce oxaliplatin on progression if neuropathy allows.
- **Second line:** switch the chemotherapy backbone (FOLFOX ↔ FOLFIRI). After first-line bevacizumab, continue an antiangiogenic agent (bevacizumab preferred over aflibercept); after FOLFOX + anti-EGFR, give FOLFIRI + bevacizumab (and vice versa). Combine chemotherapy with an antiangiogenic or an anti-EGFR antibody, never both. Patients who cannot tolerate intensive therapy can have sequential single agents, capecitabine + bevacizumab, or supportive care alone.
- **Later lines:** trifluridine–tipiracil with bevacizumab (median survival 10.8 vs 7.5 months over trifluridine–tipiracil alone; [SUNLIGHT](#)^[52], after [RECOURSE](#)^[53]); regorafenib ([CORRECT](#)^[54]); fruquintinib ([FRESKO-2](#)^[55]). Gains are measured in weeks to months: weigh them with the patient against toxicity and cost.
- **HER2-overexpressing** (pMMR, RAS wild-type) after chemotherapy: tucatinib + trastuzumab preferred ([MOUNTAINEER](#)^[13]); trastuzumab with lapatinib or pertuzumab if unavailable; trastuzumab deruxtecan later ([DESTINY-CRC01](#)^[14]; [DESTINY-CRC02](#)^[56]), and as the second-line choice for HER2-positive RAS-mutant disease. **KRAS G12C** after chemotherapy and anti-VEGF: sotorasib + panitumumab ([CodeBreak 300](#)^[10]; [survival analysis](#)^[57]) or adagrasib + cetuximab ([KRYSTAL-1](#)^[58]). **NTRK fusion:** larotrectinib, entrectinib or repotrectinib. **Anti-EGFR rechallenge** works mainly when plasma ctDNA shows RAS wild-type again after a break; rechallenge with earlier regimens is also an option when everything has been used.
- **Older and less fit patients:** decide on function and comorbidity, not age (geriatric assessment; CARG or CRASH toxicity scores; dose renally cleared drugs by creatinine clearance). Fit older adults get the same treatment as younger ones, often a doublet without the 5-FU bolus; avoid bevacizumab within 6–12 months of a stroke, myocardial infarction or arterial thrombosis. Less fit: reduced-intensity doublet, fluoropyrimidine ± bevacizumab, or trifluridine–tipiracil + bevacizumab; dMMR → single-agent immunotherapy. ECOG 2: treat selected patients, especially when the decline is due to the cancer; ECOG 3–4: supportive care for most.
- **In India:** biosimilar bevacizumab and generic capecitabine, oxaliplatin and irinotecan make doublet chemotherapy with bevacizumab widely affordable; anti-EGFR antibodies, immunotherapy and BRAF/HER2 drugs often need assistance programmes. Chemotherapy alone remains a valid, evidence-based choice when biologicals are unaffordable.




Figure 11. Side effects by drug (schematic). Signature toxicities guide counselling: cold-triggered neuropathy with oxaliplatin, rash and low magnesium with anti-EGFR antibodies, hypertension and bleeding with anti-VEGF drugs.

Quick-reference tables (also printable as one PDF from the Printable Teaching Resources page). No doses; check current approvals and product information.

Table A. Biomarkers and matched therapy

Biomarker	Frequency	Test	What it changes	Key trials	Watch for
dMMR / MSI-high	~15% of stage II, ~4% of metastatic	IHC for MLH1, MSH2, MSH6, PMS2 (or PCR/NGS MSI) on every new cancer	Stage II: no fluoropyrimidine alone. Stage III: FOLFOX + atezolizumab. Localised: neoadjuvant IO (organ sparing in rectum). Metastatic: nivolumab + ipilimumab preferred; single-agent PD-1 if less fit	KEYNOTE-177 ^[50] , CheckMate 8HW ^[48] , NICHE-2 ^[42] , Cercek 2022 ^[59] , ATOMIC ^[36]	Lynch work-up; immune side effects; pseudoprogression early
RAS (KRAS/NRAS exons 2, 3, 4)	~45–50%	Extended RAS by NGS or PCR before any anti-EGFR	Mutant → no cetuximab or panitumumab	PRIME ^[9] , CRYSTAL ^[45]	Do not use KRAS exon 2 alone
KRAS G12C	~3%	NGS	Sotorasib + panitumumab (or adagrasib + cetuximab) after chemotherapy and anti-VEGF	CodeBreaK 300 ^[10] , KRYSTAL-1 ^[58]	Single-agent KRAS inhibitors work poorly in colorectal cancer

Biomarker	Frequency	Test	What it changes	Key trials	Watch for
BRAF V600E	~8–10%	NGS or PCR; (also used to sort MLH1 loss)	First line (any side): encorafenib + cetuximab + FOLFOX (FOLFIRI alternative); later: encorafenib + cetuximab	BREAKWATER ^[11] , BEACON ^[12]	Rapid course; arthralgia, skin papillomas, fatigue
HER2 amplification	~3–5% (RAS wild-type, left)	IHC 3+, or 2+ with ISH; NGS copy number	After chemotherapy: tucatinib + trastuzumab preferred; trastuzumab deruxtecan later (or second line if RAS-mutant)	MOUNTAINEER ^[13] , DESTINY-CRCO1 ^[14] , DESTINY-CRCO2 ^[56]	Predicts anti-EGFR resistance; ILD with deruxtecan
POLE / POLD1 proofreading mutation	About 1%; ultramutated (TMB often >50/Mb)	NGS (somatic); germline testing if suspected PPAP	Metastatic: immune checkpoint inhibitor first line, as for dMMR	Guideline-based	Germline PPAP: colonoscopy from 25–30
PI3K pathway (PIK3CA, PIK3R1, PTEN)	About a third	NGS on the resected tumour	Stage I–III after surgery: low-dose aspirin for 3 years	ALASCCA ^[15]	Bleeding risk; not shown in unselected patients
NTRK, RET fusions	<1% (mostly MSI-high, right)	RNA NGS	Larotrectinib or entrectinib; selpercatinib (tumour-agnostic)	Basket trials	Rare: test if pan-wild-type
DPYD variants	A few per cent	Germline genotype (or phenotype) before fluoropyrimidines	Reduce or avoid 5-FU/capecitabine	Guidance (EMA, CPIC)	Severe mucositis, diarrhoea, neutropenia
UGT1A1*28	Common in South Asians (heterozygous)	Germline genotype	Lower irinotecan exposure in homozygotes	Guidance	Neutropenia, diarrhoea

Table B. Systemic therapy by setting (indications only; no doses)

Setting	Preferred options	Alternatives / when to modify	Evidence
Stage II high-risk colon	CAPOX 3 months, or FOLFOX 6 months	Fluoropyrimidine alone if over 70 or unfit for oxaliplatin; none if dMMR	MOSAIC ^[30] , QUASAR ^[35] , IDEA ^[34]
Stage III colon, low risk (T1–3 N1)	CAPOX 3 months	FOLFOX 6 months; capecitabine or 5-FU alone if oxaliplatin unsuitable	IDEA ^[34] , X-ACT ^[33]
Stage III colon, high risk (T4 or N2)	CAPOX or FOLFOX 6 months	CAPOX 3 months if neuropathy or poor tolerance	IDEA ^[34] , MOSAIC OS ^[31]
Stage III colon, dMMR	FOLFOX + atezolizumab	Standard FOLFOX/CAPOX where atezolizumab unavailable	ATOMIC ^[36]
Stage I–III, PI3K-pathway altered	Low-dose aspirin for 3 years after surgery (with or after adjuvant therapy)	Unselected patients: discuss; trials negative	ALASCCA ^[15] , ASCOLT ^[37]

Setting	Preferred options	Alternatives / when to modify	Evidence
Locally advanced colon (radiological T4, bulky T3)	Neoadjuvant FOLFOX (6 weeks) then surgery	Neoadjuvant nivolumab + ipilimumab if dMMR (trial or expert setting)	FOX TROT^[41] , NICHE-2^[42]
Locally advanced rectum (pMMR)	Total neoadjuvant therapy: short-course RT → chemo, or CRT + consolidation chemo, or induction chemo → CRT	Selected cT2N1/cT3N0–1 with clear MRF: FOLFOX first, RT only if poor response	RAPIDO^[60] , OPRA^[61] , PRODIGE 23^[62] , PROSPECT^[63]
Locally advanced rectum, dMMR	Anti-PD-1 alone (dostarlimab) with close MRI/endoscopy follow-up	Chemoradiation if IO unavailable (response to chemotherapy is poorer)	Cercek 2022^[59]
Metastatic, first line, pMMR, RAS/BRAF wild-type, left	FOLFOX or FOLFIRI + panitumumab or cetuximab	Bevacizumab instead if anti-EGFR unaffordable or poorly tolerated	PARADIGM^[16] , FIRE-3^[43] , CALGB/SWOG 80405^[44]
Metastatic, first line, right-sided or RAS-mutant	FOLFOX / CAPOX / FOLFIRI + bevacizumab; FOLFOXIRI + bevacizumab if fit with high burden	Right, wild-type, bevacizumab contraindicated: chemotherapy alone (not anti-EGFR); FOLFIRI if oxaliplatin within 12 months; frail: capecitabine ± bevacizumab	Hurwitz^[46] , TRIBE^[47] , sidedness analysis^[17]
Metastatic, first line, dMMR / MSI-high or POLE-ultramutated	Nivolumab + ipilimumab	Pembrolizumab, nivolumab or dostarlimab alone if less fit; 1–2 years	CheckMate 8HW^[48] , KEYNOTE-177^[50]
Metastatic, first line, BRAF V600E	Encorafenib + cetuximab + FOLFOX	FOLFIRI backbone if neuropathy	BREAKWATER^[11]
Liver-only, potentially resectable (conversion)	RAS/BRAF wild-type: oxaliplatin doublet/triplet (± anti-EGFR if left); RAS/BRAF-mutant: FOLFOXIRI + bevacizumab	≤4 cycles before surgery; about 6 months in total	CAIRO5^[23] , EPOC^[21]
Maintenance	Fluoropyrimidine + bevacizumab after 4–6 months of induction; stop oxaliplatin at 3–4 months	Treatment break with close scans for selected patients	CAIRO3^[51]
Second line	Switch backbone (FOLFOX ↔ FOLFIRI) ± continue bevacizumab, or aflibercept / ramucirumab with FOLFIRI	Encorafenib + cetuximab (BRAF); KRAS G12C or HER2 options	BEACON^[12]
Older or less fit	Doublet without 5-FU bolus if fit; otherwise fluoropyrimidine ± bevacizumab or trifluridine–tipiracil + bevacizumab	ECOG 3–4: supportive care; avoid bevacizumab within 6–12 months of arterial events	Geriatric assessment
Third line and later	Trifluridine–tipiracil + bevacizumab	Regorafenib; fruquintinib; anti-EGFR rechallenge if ctDNA RAS wild-type	SUNLIGHT^[52] , CORRECT^[54] , FRESCO-2^[55] , RECOURSE^[53]

Table C. Neoadjuvant strategies for rectal cancer compared

Strategy	Best suited to	Main result	Caveat
Short-course RT (5 × 5 Gy) → surgery	Elderly or frail; when chemotherapy is not planned	Halved local recurrence vs surgery alone (Dutch TME^[64])	Little downstaging if surgery is immediate; delay 4–8 weeks lowers complications (Stockholm III^[65])

Strategy	Best suited to	Main result	Caveat
Long-course chemoradiation → surgery → adjuvant	Traditional standard	Preoperative better than postoperative for local control and toxicity (German CAO/ARO/AIO-94 ^[66])	No survival gain; adjuvant chemotherapy often not completed
TNT: short-course RT → 18 weeks chemo → TME	High risk: T4, N2, threatened CRM, EMVI, lateral nodes	Fewer disease-related treatment failures; pathological complete response doubled (RAPIDO ^[60])	Higher local recurrence at 5 years than CRT in the trial: careful surgery and follow-up
TNT: induction FOLFIRINOX → CRT → TME	Fit patients, high distant risk	Better disease-free and overall survival (PRODIGE 23 ^[62])	Intensive; needs good performance status
TNT: CRT → consolidation chemo → watch-and-wait if cCR	Low tumours where keeping the rectum matters	About half keep the rectum at 3–5 years (OPRA ^[61])	Regrowth in about a third of watch-and-wait patients; strict surveillance
FOLFOX → selective CRT → TME	cT2N1, cT3N0–1, MRF clear, suitable for sphincter-sparing surgery	Non-inferior disease-free survival; most avoid pelvic RT (PROSPECT ^[63])	Not for T4, threatened margin or low tumours
Anti-PD-1 alone	dMMR / MSI-high rectal cancer	Clinical complete response in nearly all patients in the first series (Cercek 2022 ^[59])	Small numbers; confirm MMR status; surveillance as for watch-and-wait

Table D. Managing common side effects (principles; follow local protocols)

Problem	Usual cause	Do	Hold or stop when
Cold-triggered tingling, cramps; later persistent numbness	Oxaliplatin	Warn about cold drinks and metal; check function every cycle; stop-and-go strategy	Persistent numbness affecting function (grade 2 lasting, or 3): stop oxaliplatin, continue the rest
Diarrhoea	Irinotecan, fluoropyrimidines, IO	Loperamide plan given in advance; fluids; check for neutropenia	Severe, bloody, febrile or with dehydration: admit; on IO consider colitis (stool tests, steroids)
Severe toxicity in cycle 1 (mucositis, diarrhoea, neutropenia)	DPD deficiency	Stop fluoropyrimidine; supportive care; uridine triacetate where available	Do not rechallenge at full intensity; genotype DPYD
Chest pain on infusion or capecitabine	Fluoropyrimidine coronary vasospasm	Stop, ECG and troponin	Recurrence is common: switch to raltitrexed or non-fluoropyrimidine regimen
Painful red palms and soles	Capecitabine; regorafenib (hand-foot skin reaction)	Emollients, urea cream, avoid friction; dose delay	Blistering or limits self-care
Acneiform rash, paronychia	Cetuximab, panitumumab	Pre-emptive doxycycline/minocycline, moisturiser, sunscreen; rash often tracks with benefit	Severe or infected rash; hold then resume at lower intensity
Low magnesium, cramps, arrhythmia	Anti-EGFR antibodies	Check magnesium, calcium, potassium regularly; replace	Symptomatic or refractory hypomagnesaemia
Hypertension, proteinuria	Bevacizumab, regorafenib, fruquintinib	BP every visit; urine dipstick	Uncontrolled BP, nephrotic-range proteinuria

Problem	Usual cause	Do	Hold or stop when
Bleeding, perforation, wound problems	Anti-VEGF drugs	Stop 4–6 weeks before elective surgery; restart after wound heals	Perforation, major bleeding, arterial thrombosis: stop permanently; avoid with an uncovered stent
Neutropenia	Irinotecan, trifluridine–tipiracil, FOLFOXIRI	Count before each cycle; G-CSF per risk	Febrile neutropenia: emergency antibiotics
Colitis, hepatitis, thyroid, adrenal problems	Pembrolizumab, nivolumab ± ipilimumab	Baseline and periodic TSH, cortisol, liver tests; report diarrhoea early	Grade ≥2 colitis or hepatitis: hold, steroids; grade 4: stop


07 Management algorithms

These decision paths pull together sections 02–06. Colours: teal for early or favourable paths, purple for intermediate, rose for high-risk or later-line paths.

ALGORITHM 1

Non-metastatic colon cancer

Colon cancer on biopsy: complete colonoscopy · CT chest–abdomen–pelvis · CEA · MMR immunohistochemistry · fitness and nutrition · tumour board



All patients: Lynch work-up if dMMR without BRAF V600E or MLH1 methylation (Lynch: consider total colectomy with ileorectal anastomosis) · start adjuvant therapy within about 8 weeks · DPYD testing where available · PI3K-pathway alteration on NGS → low-dose aspirin for 3 years · replace vitamin D if below 20 ng/mL · colonoscopy at 1 year · exercise and weight advice from the first visit.

ALGORITHM 2

Metastatic colorectal cancer

New stage IV: CT ± MRI liver (PET only before metastasectomy) · **MMR, RAS, BRAF, HER2** (NGS) · DPYD · sidedness · fitness and goals · hepatobiliary/thoracic surgeon at the first MDT

RESECTABLE OR POTENTIALLY RESECTABLE

Low risk (≤ 4 lesions in one lobe, RAS/BRAF wild-type) → surgery first; ablation for ≤ 3 cm

Higher risk (> 4 , bilobar, RAS/BRAF-mutant) → chemotherapy first (≤ 4 cycles)

Unresectable liver-only → conversion: oxaliplatin doublet/triplet ($\pm$ anti-EGFR if left, wild-type); FOLFOXIRI + bev if RAS/BRAF-mutant

Re-image every 6–8 weeks → operate as soon as resectable; about 6 months of treatment in total

Fiducials for small, deep lesions before chemotherapy · asymptomatic primary: no upfront resection

UNRESECTABLE: FIRST LINE

dMMR or POLE-ultramutated → nivolumab + ipilimumab (single-agent anti-PD-1 if less fit)

BRAF V600E → encorafenib + cetuximab + FOLFOX (any side)

Left, RAS/BRAF wild-type → doublet + anti-EGFR

Right or RAS-mutant → doublet (triplet if fit, high burden) + bevacizumab; FOLFIRI if oxaliplatin within 12 months

Frail → capecitabine $\pm$ bevacizumab; biologicals unaffordable → chemotherapy alone

Maintenance after 4–6 months; local therapy for oligoprogression

LATER LINES

Second line: switch backbone $\pm$ anti-VEGF (never with anti-EGFR); after IO → chemotherapy; BRAF → encorafenib + cetuximab; HER2 → tucatinib + trastuzumab (T-DXd later, or second line if RAS-mutant); G12C → sotorasib + panitumumab

Third line: trifluridine–tipiracil + bevacizumab, regorafenib, fruquintinib; anti-EGFR rechallenge if ctDNA RAS wild-type

Early palliative care; clinical trials at every line

All patients: discuss goals early · obstruction → stent or diversion (avoid stents with bevacizumab) · peritoneal disease → expert-centre opinion on cytoreduction · selected unresectable liver-only disease → transplant evaluation in specialised programmes.

o8 Survivorship, follow-up & reducing recurrence




Figure 12. When relapses happen (illustrative shape, not to scale). Intensive CEA and CT surveillance is concentrated in the first 3 years because that is when a resectable relapse can still be found; colonoscopy continues for new polyps and cancers.




Figure 13. A simple follow-up rhythm (NCCN, ESMO, ASCO). More intensive testing than this did not improve survival in the FACS and COLOFOL trials, but surveillance still finds resectable relapses.

Follow-up protocol (NCCN, ESMO, ASCO)

What	How often	Notes
History, examination, CEA	Every 3–6 months for 3 years, then every 6 months in years 4–5 (CEA at least for the first 2–3 years)	Stage II–III and after resection of metastases, if the patient would be fit for further curative treatment; rectal examination after low anterior resection or local excision. A confirmed rising CEA triggers imaging.
CT chest and abdomen (pelvis for rectal cancer)	Every 6–12 months up to 5 years (NCCN), or at least yearly for 3 years	Stage I: CT not routine (except after local or endoscopic excision alone). Earlier relapses are more often resectable.
Colonoscopy	At 1 year (or 3–6 months if not complete before surgery); if an advanced adenoma is found, again at 1 year, otherwise every 3–5 years	Every 1–2 years in Lynch syndrome; yearly in PSC-associated colitis.

What	How often	Notes
Rectal endoscopy and MRI pelvis (rectum)	Watch-and-wait or local excision: every 3–4 months for 2 years, then 6-monthly to year 5. Low anterior resection without pelvic RT: proctosigmoidoscopy with endorectal ultrasound or MRI every 6 months for up to 5 years	Regrowth is usually luminal and salvageable when found early.
Not routine	—	PET-CT, other tumour markers, ctDNA outside trials.

- **Intensity:** more intensive CEA and CT schedules found more resectable relapses but did not improve overall survival in [FACS](#)^[67] and [COLOFOL](#)^[68]; the schedule above is a balance, not a minimum.
- **Exercise:** a structured 3-year programme of aerobic activity after adjuvant chemotherapy reduced recurrence or new cancer (5-year disease-free survival 80% vs 74%) and improved overall survival ([CHALLENGE](#)^[69]). Prescribe it like a drug: about 150 minutes a week of brisk walking, built up with support.
- **Other risk factors:** healthy weight (severe obesity, BMI over 35, is linked with worse outcomes); in observational studies of stage III colon cancer, a Western-style diet (red and processed meat, sweets, refined grains) and a high dietary glycaemic load were linked with more recurrence, and a prudent diet (vegetables, fruit, whole grains, fish, pulses) with less; no smoking, little or no alcohol, a plant-rich diet with less red and processed meat. Aspirin: see section 06 (PI3K-altered cancers, Lynch syndrome). Check and replace vitamin D if low.
- **Late effects:** oxaliplatin neuropathy (may persist; about 1% have chronic severe painful neuropathy needing duloxetine-type drugs or other analgesics), fatigue, depression, work and money worries, higher rates of diabetes and cardiovascular disease, radiation proctitis and pelvic fractures, low anterior resection syndrome, stoma care, urinary and sexual dysfunction, infertility and early menopause after pelvic RT, insufficiency fractures, anxiety about recurrence.
- **Stoma:** stoma nurse before and after surgery; hernia and dehydration (high-output ileostomy) are common problems; most people return to work, travel and religious practice.
- **Family:** first-degree relatives start colonoscopy at 40 (or 10 years before the patient's age at diagnosis); cascade genetic testing in Lynch and polyposis families.
- **Shared care:** give the family doctor the stage, the follow-up schedule and the red flags (below).

SHARE WITH PATIENTS: LIFE AFTER TREATMENT

Plain-language myth pages written for patients and families. Send them from the clinic to reinforce what you discussed.

[Telugu-English handout: living with a stoma \(తెలుగు-English\)](#)

[Telugu-English handout: stool test \(FIT\) and colonoscopy \(తెలుగు-English\)](#)

[Once colorectal cancer treatment is finished, follow-up isn't needed?](#)

[Every rectal cancer patient needs a permanent colostomy bag?](#)

[Living with a stoma \(bag\) means a normal life is over?](#)

[After bowel cancer treatment, rest is best and exercise should be avoided?](#)

[When one person in the family has bowel cancer, the others do not need any tests?](#)

09 Landmark trials




Figure 14. How to read a survival curve (illustrative). Use the absolute gap at a fixed time and overall survival, not the median or PFS alone, when weighing benefit against toxicity and cost.

- [National Polyp Study](#)^[6] and [NordICC](#)^[7]: does finding and removing polyps prevent death? Polypectomy roughly halved colorectal cancer deaths; a population invitation to one colonoscopy cut incidence by about a fifth, limited by low uptake. → Screening works only if people attend and positive tests are followed up.
- [MOSAIC](#)^[30] and [IDEA](#)^[34]: oxaliplatin added to fluoropyrimidine adjuvant therapy improved survival in stage III; three months of CAPOX is enough for low-risk stage III. → Match duration to risk and spare nerves.
- [FOX-TROT](#)^[41] and [NICHE-2](#)^[42]: treatment before colon surgery. Six weeks of FOLFOX downstaged locally advanced pMMR cancers; two cycles of immunotherapy eradicated most dMMR cancers. → Test MMR on the biopsy, before surgery.
-
-
-
- [KEYNOTE-177](#)^[50] and [CheckMate 8HW](#)^[48]: immunotherapy beats chemotherapy for metastatic MSI-high cancer, with long-lasting responses. → MMR testing at diagnosis of metastatic disease is mandatory.
- [PARADIGM](#)^[16]: panitumumab versus bevacizumab with FOLFOX in RAS wild-type disease. Overall survival improved in left-sided tumours (median about 38 vs 34 months). → Sidedness and RAS together choose the biological.
- [BREAKWATER](#)^[11]: first-line encorafenib + cetuximab + FOLFOX for BRAF V600E roughly doubled median survival. → BRAF testing belongs at diagnosis.
- [CHALLENGE](#)^[69]: a supported exercise programme after adjuvant chemotherapy reduced recurrence and death. → Exercise is part of treatment.
- [ALASCCA](#)^[15]: low-dose aspirin for 3 years after surgery for PI3K-altered stage I–III colorectal cancer roughly halved recurrence, whereas trials in unselected patients were negative. → A cheap drug, selected by a

tumour biomarker.

- [COLLISION](#)^[24]: thermal ablation versus resection for small liver metastases (3 cm or less). Survival was similar with fewer complications. → Ablation is a curative-intent option, chosen in the MDT.

10 Practice pearls & pitfalls

- Piles do not explain anaemia. Iron-deficiency anaemia in a man or a post-menopausal woman means colonoscopy and upper endoscopy, not iron tablets alone.
- Do a digital rectal examination in every patient with rectal bleeding; most low rectal cancers are within reach of a finger.
- A negative FIT does not override red-flag symptoms, and CEA never rules cancer in or out.
- Take multiple biopsies of any ileocaecal mass for histology and TB tests; do not start antituberculous treatment without proof.
- Request MMR status on the first biopsy. It decides neoadjuvant immunotherapy, adjuvant chemotherapy in stage II and screening of the family.
-
-
-
- Fewer than 12 nodes in the specimen is a quality problem and a high-risk feature, not reassurance.
- Start adjuvant chemotherapy within about 8 weeks; three months of CAPOX is enough for low-risk stage III.
- Test DPYD where available, and stop a fluoropyrimidine at once after severe cycle-1 toxicity.
- Never give anti-EGFR antibodies without an extended RAS (and BRAF) result; prefer them for left-sided tumours.
- Show every patient with liver or lung metastases to a hepatobiliary or thoracic surgeon before chemotherapy starts.
- Chemotherapy with biosimilar bevacizumab is an evidence-based, affordable first-line option; do not let unaffordable biologicals delay treatment.
- Send the resected tumour for NGS where affordable: a PI3K-pathway alteration means 3 years of low-dose aspirin (ALASCCA), and POLE/POLD1 or dMMR results change metastatic treatment.
- Patients with primary sclerosing cholangitis need colonoscopy at diagnosis and yearly if they have colitis; right-sided cancers in their 40s are typical.
- A positive germline result is rarely an emergency; a variant of uncertain significance changes nothing. Single MUTYH carriers screen like everyone else.
- Prescribe exercise, check neuropathy at every visit, and arrange colonoscopy for first-degree relatives.

11 Case vignettes: lines of management

Four cases, written for postgraduates and practising clinicians. Read the presentation, decide on stage, workup and plan, then open the line of management. No doses are given; durations follow the evidence.

CASE 1 Colon · stage III · dMMR · Lynch

Six months of iron tablets in a 34-year-old

34 years, woman

pT3 N1b (stage IIIB)

MSH2 and MSH6 lost

A 34-year-old teacher has had fatigue for a year and was treated with iron tablets twice for haemoglobin of 8 g/dL with low ferritin. Her stools are normal. Her father had “intestinal cancer” at 48 and her aunt had uterine cancer at 50.

Before you open the answer: what is the stage, what must be done before treatment, and what is the plan?

1 INVESTIGATE THE ANAEMIA

Iron deficiency without an obvious cause needs both endoscopies, even at 34. Colonoscopy: ulcerated caecal cancer; biopsy adenocarcinoma, poorly differentiated with tumour-infiltrating lymphocytes.

2 STAGE AND TEST

CT chest–abdomen–pelvis: no metastases. CEA normal. MMR immunohistochemistry: loss of MSH2 and MSH6 — Lynch syndrome likely.

3 SURGERY

Lynch syndrome is suspected before surgery, so total colectomy with ileorectal anastomosis is offered to cut the roughly 20% 10-year risk of a new colon cancer after segmental resection; she chooses it after discussing bowel function. 31 nodes, 3 positive (pT3 N1b).

4 ADJUVANT THERAPY

Stage III dMMR: FOLFOX with atezolizumab for 6 months where available (ATOMIC); otherwise CAPOX or FOLFOX.

5 GERMLINE AND FAMILY

Germline MSH2 pathogenic variant confirmed. Yearly endoscopy of the remaining rectum; upper endoscopy from 30–35 with H. pylori treatment; yearly urinalysis (MSH2 carries urothelial risk); risk-reducing hysterectomy and salpingo-oophorectomy after childbearing; daily aspirin. Cascade testing offered to siblings and children (from 18–20).

TEACHING POINTS

- Anaemia in a young adult with low ferritin is a gut question, not a dietary one.
- MMR status changes adjuvant therapy, the extent of surgery and the family's screening.
- MSH2/MSH6 loss is not explained by MLH1 methylation: go straight to germline testing.

See: Sections 02, 04, 06 · Algorithm 1 · [ATOMIC](#)^[36], [CAPP2](#)^[2], [Parry 2011](#)^[70]

CASE 2 Colon · obstruction · stage II

A distended abdomen in a 72-year-old farmer

72 years, diabetic

pT4a NO (stage IIB)

pMMR

A 72-year-old man presents with 4 days of absolute constipation, colicky pain and a distended abdomen. He was treated for “gas” and “piles” over the past 3 months.

Before you open the answer: what is the stage, what must be done before treatment, and what is the plan?

1 RESUSCITATE AND IMAGE

Drip and suck; contrast CT: obstructing sigmoid tumour, dilated colon with a caecum of 9 cm, no perforation, no metastases.

2 CHOOSE THE OPERATION

Options: emergency resection (Hartmann's, or primary anastomosis ± covering stoma), or a self-expanding stent as a bridge to elective laparoscopic surgery in 1–2 weeks in an experienced centre. He had a stent, recovered, completed staging and colonoscopy, and had an elective sigmoid colectomy.

3 PATHOLOGY

pT4a N0, 16 nodes, pMMR, no lymphovascular invasion. Obstruction and T4 are both high-risk features.

4 ADJUVANT THERAPY

High-risk stage II: 3 months of CAPOX, or a fluoropyrimidine alone given his age and neuropathy from diabetes; absolute benefit is modest and is discussed with him.

5 FOLLOW-UP

CEA and CT for 5 years; colonoscopy at 1 year; exercise programme.

TEACHING POINTS

- Emergency presentation is common and carries worse survival; earlier colonoscopy would have helped.
- Stent as a bridge allows staging and elective surgery but needs expertise.
- Stage II benefit from chemotherapy is small: risk features and fitness decide.

See: Sections 02, 03, 05, 06 · Algorithm 1 · [IDEA](#)^[34], [QUASAR](#)^[35]

CASE 3 Metastatic · liver-only · conversion**Six liver metastases in a 55-year-old**

55 years, man

Sigmoid primary, 6 liver lesions

RAS/BRAF wild-type, pMMR, HER2-negative

A 55-year-old accountant has a sigmoid cancer with six liver metastases in both lobes, the largest 6 cm near the right hepatic vein. CT chest is clear. He is fit (ECOG 0).

Before you open the answer: what is the stage, what must be done before treatment, and what is the plan?

1 MDT WITH A LIVER SURGEON

Not resectable now: the remnant would be too small. Potentially resectable if the tumours shrink.

2 CONVERSION THERAPY

Left-sided, RAS/BRAF wild-type: FOLFOX (or FOLFIRI) with panitumumab or cetuximab for the highest response rate.

3 RESTAGE EVERY 2–3 MONTHS

After 4 months: 60% shrinkage; now resectable with a two-stage approach (left lateral clearance, portal vein embolisation, then right hepatectomy).

4 SURGERY

Liver resections and sigmoid colectomy in staged procedures; RO. Ablation for a small residual lesion.

5 AFTER

Complete about 6 months of perioperative chemotherapy in total; CEA and CT every 3–4 months; about 4 in 10 such patients are alive at 5 years, though most relapse at some point.

TEACHING POINTS

- Show liver metastases to a hepatobiliary surgeon before chemotherapy.
- Choose the most active regimen for the biomarkers when conversion is the goal.
- Operate as soon as resectable: “vanishing” lesions and chemotherapy liver injury complicate late surgery.

See: Sections 05, 06 · Algorithm 3 · [PARADIGM](#)^[16], [EPOC](#)^[21]

CASE 4 Metastatic · BRAF V600E**Ascites and right-sided cancer in a 47-year-old**

47 years, woman

Ascending colon, peritoneal + liver

BRAF V600E, pMMR

A 47-year-old homemaker presents with abdominal distension and weight loss. CT shows an ascending colon mass, omental cake, ascites and three liver metastases. Biopsy: adenocarcinoma.

Before you open the answer: what is the stage, what must be done before treatment, and what is the plan?

1**BIOMARKERS FIRST**

NGS: BRAF V600E, RAS wild-type, pMMR, HER2-negative. DPYD normal. ECOG 1.

2**FIRST LINE**

Encorafenib + cetuximab + mFOLFOX6 (BREAKWATER) where accessible; otherwise FOLFOXIRI + bevacizumab if fit, or FOLFOX + bevacizumab. Anti-EGFR alone with chemotherapy is ineffective in BRAF-mutant disease.

3**SUPPORTIVE**

Early palliative care; nutrition; drainage of ascites as needed; watch for bowel obstruction from peritoneal disease.

4**SECOND LINE**

If not given first: encorafenib + cetuximab (BEACON). Later: trifluridine–tipiracil + bevacizumab, regorafenib or fruquintinib; clinical trials.

5**GOALS**

Discuss prognosis honestly: BRAF V600E disease progresses fast, but targeted combinations have substantially improved survival.

TEACHING POINTS

- Test BRAF at diagnosis of metastatic disease: it changes first-line therapy.
- Right-sided, peritoneal and BRAF often go together.
- Plan supportive and palliative care from the start.

See: Sections 04, 06 · Algorithm 3 · [BREAKWATER](#)^[11], [BEACON](#)^[12]

12 References & guidelines

- NCCN Clinical Practice Guidelines in Oncology: Colon Cancer; Rectal Cancer; Colorectal Cancer Screening; Genetic/Familial High-Risk Assessment: Colorectal, Endometrial and Gastric (current versions).
- ESMO Clinical Practice Guidelines: localised colon cancer; rectal cancer; metastatic colorectal cancer (and ESMO Magnitude of Clinical Benefit Scale).
- ASCO guidelines: treatment of metastatic colorectal cancer; surveillance after curative treatment; management of locally advanced rectal cancer.
- AJCC/UICC TNM Classification, 8th edition; College of American Pathologists colorectal reporting protocol; ESGAR rectal MRI consensus.
- US Multi-Society Task Force and USPSTF 2021 screening recommendations; British Society of Gastroenterology / NICE DG30 guidance on FIT in symptomatic patients; ESGE guidance on colonic stenting.
- ICMR consensus document for the management of colorectal cancer; Indian Council of Medical Research National Cancer Registry Programme reports; GLOBOCAN 2022.
- Cross-checked in October 2026 against current evidence summaries (UpToDate topic reviews on epidemiology and risk factors, molecular genetics, polyps, screening strategies and tests, family-history screening, Lynch syndrome, MUTYH and POLE/POLD1 gene-test interpretation, PSC-associated colitis, presentation and staging, pathology and prognosis, management of primary colon cancer, adjunctive aspirin and vitamin D, survivorship, diet and exercise, post-treatment surveillance, systemic therapy principles, initial and later-line therapy for metastatic disease, older adults, liver resection and potentially resectable liver metastases, non-surgical liver-directed therapy and locoregional management). All text is written in our own words and cites primary sources; regimen dose tables were not used.
- Trial names above link to PubMed searches for the primary publications.

Core trials cited on this page (70)

Journal, year, volume and first page; each links to the PubMed record by exact citation .

NUMBERED IN ORDER OF FIRST CITATION IN THE TEXT

1. **PSC-IBD colorectal neoplasia meta-analysis:** Zheng HH, Jiang XL. Increased risk of colorectal neoplasia in patients with primary sclerosing cholangitis and inflammatory bowel disease: a meta-analysis of 16 observational studies. *Eur J Gastroenterol Hepatol* 2016;28:383. [PubMed](#) [Screening and prevention](#)
2. **CAPP2 long-term:** Burn J, Sheth H, Elliott F, et al. Cancer prevention with aspirin in hereditary colorectal cancer (Lynch syndrome), 10-year follow-up and registry-based 20-year data in the CAPP2 study. *Lancet* 2020;395:1855. [PubMed](#) [Screening and prevention](#)
3. **MUTYH carrier risk (Win 2014):** Win AK, Dowty JG, Cleary SP, et al. Risk of colorectal cancer for carriers of mutations in MUTYH, with and without a family history of cancer. *Gastroenterology* 2014;146:1208. [PubMed](#) [Screening and prevention](#)
4. **PPAP clinical features (Palles 2022):** Palles C, Martin L, Domingo E, et al. The clinical features of polymerase proof-reading associated polyposis (PPAP) and recommendations for patient management. *Fam Cancer* 2022;21:197. [PubMed](#) [Screening and prevention](#)
5. **Minnesota FOBT trial:** Mandel JS, Bond JH, Church TR, et al. Reducing mortality from colorectal cancer by screening for fecal occult blood. *N Engl J Med* 1993;328:1365. [PubMed](#) [Screening and prevention](#)
6. **National Polyp Study:** Zauber AG, Winawer SJ, O'Brien MJ, et al. Colonoscopic polypectomy and long-term prevention of colorectal-cancer deaths. *N Engl J Med* 2012;366:687. [PubMed](#) [Screening and prevention](#)
7. **NordICC:** Bretthauer M, Løberg M, Wieszczyn P, et al. Effect of colonoscopy screening on risks of colorectal cancer and related death. *N Engl J Med* 2022;387:1547. [PubMed](#) [Screening and prevention](#)
8. **USPSTF 2021 colorectal:** US Preventive Services Task Force. Screening for colorectal cancer: US Preventive Services Task Force recommendation statement. *JAMA* 2021;325:1965. [PubMed](#) [Screening and prevention](#)

9. **PRIME:** Douillard JY, Oliner KS, Siena S, et al. Panitumumab–FOLFOX4 treatment and RAS mutations in colorectal cancer. *N Engl J Med* 2013;369:1023. [PubMed](#) Metastatic disease
10. **CodeBreak 300:** Fakih MG, Salvatore L, Esaki T, et al. Sotorasib plus panitumumab in refractory colorectal cancer with mutated KRAS G12C. *N Engl J Med* 2023;389:2125. [PubMed](#) Metastatic disease
11. **BREAKWATER:** Elez E, Yoshino T, Shen L, et al. Encorafenib, cetuximab, and mFOLFOX6 in BRAF–mutated colorectal cancer. *N Engl J Med* 2025;392:2425. [PubMed](#) Metastatic disease
12. **BEACON:** Kopetz S, Grothey A, Yaeger R, et al. Encorafenib, binimetinib, and cetuximab in BRAF V600E–mutated colorectal cancer. *N Engl J Med* 2019;381:1632. [PubMed](#) Metastatic disease
13. **MOUNTAINEER:** Strickler JH, Cercek A, Siena S, et al. Tucatinib plus trastuzumab for chemotherapy–refractory, HER2–positive, RAS wild–type unresectable or metastatic colorectal cancer (MOUNTAINEER). *Lancet Oncol* 2023;24:496. [PubMed](#) Metastatic disease
14. **DESTINY–CRC01:** Siena S, Di Bartolomeo M, Raghav K, et al. Trastuzumab deruxtecan (DS–8201) in patients with HER2–expressing metastatic colorectal cancer (DESTINY–CRC01). *Lancet Oncol* 2021;22:779. [PubMed](#) Metastatic disease
15. **ALASCCA:** Martling A, Hed Myrberg I, Nilbert M, et al. Low–dose aspirin for PI3K–altered localized colorectal cancer. *N Engl J Med* 2025;393:1051. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
16. **PARADIGM:** Watanabe J, Muro K, Shitara K, et al. Panitumumab vs bevacizumab added to standard first–line chemotherapy and overall survival among patients with RAS wild–type, left–sided metastatic colorectal cancer. *JAMA* 2023;329:1271. [PubMed](#) Metastatic disease
17. **Sidedness pooled analysis:** Arnold D, Lueza B, Douillard JY, et al. Prognostic and predictive value of primary tumour side in patients with RAS wild–type metastatic colorectal cancer treated with chemotherapy and EGFR directed antibodies in six randomized trials. *Ann Oncol* 2017;28:1713. [PubMed](#) Metastatic disease
18. **DYNAMIC:** Tie J, Cohen JD, Lahouel K, et al. Circulating tumor DNA analysis guiding adjuvant therapy in stage II colon cancer. *N Engl J Med* 2022;386:2261. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
19. **COST:** Clinical Outcomes of Surgical Therapy Study Group. A comparison of laparoscopically assisted and open colectomy for colon cancer. *N Engl J Med* 2004;350:2050. [PubMed](#) Surgery and local therapy
20. **COLOR:** Buunen M, Veldkamp R, Hop WC, et al. Survival after laparoscopic surgery versus open surgery for colon cancer: long–term outcome of a randomised clinical trial. *Lancet Oncol* 2009;10:44. [PubMed](#) Surgery and local therapy
21. **EPOC:** Nordlinger B, Sorbye H, Glimelius B, et al. Perioperative chemotherapy with FOLFOX4 and surgery versus surgery alone for resectable liver metastases from colorectal cancer (EORTC 40983). *Lancet* 2008;371:1007. [PubMed](#) Surgery and local therapy
22. **EPOC OS:** Nordlinger B, Sorbye H, Glimelius B, et al. Perioperative FOLFOX4 chemotherapy and surgery versus surgery alone for resectable liver metastases from colorectal cancer (EORTC 40983): long–term results. *Lancet Oncol* 2013;14:1208. [PubMed](#) Surgery and local therapy
23. **CAIRO5:** Bond MJG, Bolhuis K, Loosveld OJL, et al. First–line systemic treatment strategies in patients with initially unresectable colorectal cancer liver metastases (CAIRO5): an open–label, multicentre, randomised, controlled, phase 3 study. *Lancet Oncol* 2023;24:757. [PubMed](#) Surgery and local therapy
24. **COLLISION:** van der Lei S, Puijk RS, Dijkstra M, et al. Thermal ablation versus surgical resection of small–size colorectal liver metastases (COLLISION): an international, randomised, controlled, phase 3 non–inferiority trial. *Lancet Oncol* 2025;26:187. [PubMed](#) Surgery and local therapy
25. **SIRT combined analysis (FOXFIRE/SIRFLOX):** Wasan HS, Gibbs P, Sharma NK, et al. First–line selective internal radiotherapy plus chemotherapy versus chemotherapy alone in patients with liver metastases from colorectal cancer (FOXFIRE, SIRFLOX, and FOXFIRE–Global): a combined analysis of three randomised phase 3 trials. *Lancet Oncol* 2017;18:1159. [PubMed](#) Surgery and local therapy
26. **PRODIGE 7:** Quénet F, Elias D, Roca L, et al. Cytoreductive surgery plus hyperthermic intraperitoneal chemotherapy versus cytoreductive surgery alone for colorectal peritoneal metastases (PRODIGE 7). *Lancet Oncol* 2021;22:256. [PubMed](#) Surgery and local therapy
27. **JCOG1007:** Kanemitsu Y, Shitara K, Mizusawa J, et al. Primary tumor resection plus chemotherapy versus chemotherapy alone for colorectal cancer patients with asymptomatic, synchronous unresectable metastases (JCOG1007; iPACS). *J Clin Oncol* 2021;39:1098. [PubMed](#) Surgery and local therapy
28. **SYNCHRONOUS/CCRe–IV:** Rahbari NN, Biondo S, Frago R, et al. Primary tumor resection before systemic therapy in patients with colon cancer and unresectable metastases: combined results of the SYNCHRONOUS and CCRe–IV trials. *J Clin Oncol* 2024;42:1531. [PubMed](#) Surgery and local therapy
29. **TRANSMET:** Adam R, Piedvache C, Chiche L, et al. Liver transplantation plus chemotherapy versus chemotherapy alone in patients with permanently unresectable colorectal liver metastases (TransMet): results from a multicentre, open–label, prospective, randomised controlled trial. *Lancet* 2024;404:1107. [PubMed](#) Surgery and local therapy

30. **MOSAIC:** André T, Boni C, Mounedji-Boudiaf L, et al. Oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for colon cancer. *N Engl J Med* 2004;350:2343. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
31. **MOSAIC OS:** André T, Boni C, Navarro M, et al. Improved overall survival with oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment in stage II or III colon cancer in the MOSAIC trial. *J Clin Oncol* 2009;27:3109. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
32. **NSABP C-07:** Kuebler JP, Wieand HS, O'Connell MJ, et al. Oxaliplatin combined with weekly bolus fluorouracil and leucovorin as surgical adjuvant chemotherapy for stage II and III colon cancer: results from NSABP C-07. *J Clin Oncol* 2007;25:2198. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
33. **X-ACT:** Twelves C, Wong A, Nowacki MP, et al. Capecitabine as adjuvant treatment for stage III colon cancer. *N Engl J Med* 2005;352:2696. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
34. **IDEA:** Grothey A, Sobrero AF, Shields AF, et al. Duration of adjuvant chemotherapy for stage III colon cancer. *N Engl J Med* 2018;378:1177. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
35. **QUASAR:** QUASAR Collaborative Group. Adjuvant chemotherapy versus observation in patients with colorectal cancer: a randomised study. *Lancet* 2007;370:2020. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
36. **ATOMIC:** Sinicrope FA, Ou FS, Arnold D, et al. Atezolizumab plus FOLFOX for stage III mismatch repair-deficient colon cancer. *N Engl J Med* 2026;394:1155. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
37. **ASCOLT:** Chia JWK, Segelov E, Deng Y, et al. Aspirin after completion of standard adjuvant therapy for colorectal cancer (ASCOLT): an international, multicentre, phase 3, randomised, double-blind, placebo-controlled trial. *Lancet Gastroenterol Hepatol* 2025;10:198. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
38. **CALGB/SWOG 80702:** Meyerhardt JA, Shi Q, Fuchs CS, et al. Effect of celecoxib vs placebo added to standard adjuvant therapy on disease-free survival among patients with stage III colon cancer: the CALGB/SWOG 80702 (Alliance) randomized clinical trial. *JAMA* 2021;325:1277. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
39. **CALGB/SWOG 80702 PIK3CA:** Nowak JA, Twombly T, Ma C, et al. Improved survival with adjuvant cyclooxygenase 2 inhibition in PIK3CA-activated stage III colon cancer: CALGB/SWOG 80702 (Alliance). *J Clin Oncol* 2024;42:2853. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
40. **Vitamin D supplementation meta-analysis:** Vaughan-Shaw PG, Buijs LF, Blackmur JP, et al. The effect of vitamin D supplementation on survival in patients with colorectal cancer: systematic review and meta-analysis of randomised controlled trials. *Br J Cancer* 2020;123:1705. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
41. **FOxTROT:** Morton D, Seymour M, Magill L, et al. Preoperative chemotherapy for operable colon cancer: mature results of an international randomized controlled trial. *J Clin Oncol* 2023;41:1541. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
42. **NICHE-2:** Chalabi M, Verschoor YL, Tan PB, et al. Neoadjuvant immunotherapy in locally advanced mismatch repair-deficient colon cancer. *N Engl J Med* 2024;390:1949. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
43. **FIRE-3:** Heinemann V, von Weikersthal LF, Decker T, et al. FOLFIRI plus cetuximab versus FOLFIRI plus bevacizumab as first-line treatment for patients with metastatic colorectal cancer (FIRE-3). *Lancet Oncol* 2014;15:1065. [PubMed](#) Metastatic disease
44. **CALGB/SWOG 80405:** Venook AP, Niedzwiecki D, Lenz HJ, et al. Effect of first-line chemotherapy combined with cetuximab or bevacizumab on overall survival in patients with KRAS wild-type advanced or metastatic colorectal cancer. *JAMA* 2017;317:2392. [PubMed](#) Metastatic disease
45. **CRYSTAL:** Van Cutsem E, Köhne CH, Hitre E, et al. Cetuximab and chemotherapy as initial treatment for metastatic colorectal cancer. *N Engl J Med* 2009;360:1408. [PubMed](#) Metastatic disease
46. **Hurwitz bevacizumab trial:** Hurwitz H, Fehrenbacher L, Novotny W, et al. Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med* 2004;350:2335. [PubMed](#) Metastatic disease
47. **TRIBE:** Loupakis F, Cremolini C, Masi G, et al. Initial therapy with FOLFOXIRI and bevacizumab for metastatic colorectal cancer. *N Engl J Med* 2014;371:1609. [PubMed](#) Metastatic disease
48. **CheckMate 8HW:** André T, Elez E, Van Cutsem E, et al. Nivolumab plus ipilimumab in microsatellite-instability-high metastatic colorectal cancer. *N Engl J Med* 2024;391:2014. [PubMed](#) Metastatic disease
49. **CheckMate 8HW (vs nivolumab):** André T, Elez E, Lenz HJ, et al. Nivolumab plus ipilimumab versus nivolumab in microsatellite instability-high metastatic colorectal cancer (CheckMate 8HW): a randomised, open-label, phase 3 trial. *Lancet* 2025;405:383. [PubMed](#) Metastatic disease
50. **KEYNOTE-177:** André T, Shiu KK, Kim TW, et al. Pembrolizumab in microsatellite-instability-high advanced colorectal cancer. *N Engl J Med* 2020;383:2207. [PubMed](#) Metastatic disease
51. **CAIRO3:** Simkens LH, van Tinteren H, May A, et al. Maintenance treatment with capecitabine and bevacizumab in metastatic colorectal cancer (CAIRO3). *Lancet* 2015;385:1843. [PubMed](#) Metastatic disease
52. **SUNLIGHT:** Prager GW, Taieb J, Fakih M, et al. Trifluridine-tipiracil and bevacizumab in refractory metastatic colorectal cancer. *N Engl J Med* 2023;388:1657. [PubMed](#) Metastatic disease

53. **RECURSE:** Mayer RJ, Van Cutsem E, Falcone A, et al. Randomized trial of TAS-102 for refractory metastatic colorectal cancer. *N Engl J Med* 2015;372:1909. [PubMed](#) Metastatic disease
54. **CORRECT:** Grothey A, Van Cutsem E, Sobrero A, et al. Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT). *Lancet* 2013;381:303. [PubMed](#) Metastatic disease
55. **FRESCO-2:** Dasari A, Lonardi S, Garcia-Carbonero R, et al. Fruquintinib versus placebo in patients with refractory metastatic colorectal cancer (FRESCO-2). *Lancet* 2023;402:41. [PubMed](#) Metastatic disease
56. **DESTINY-CRCO2:** Raghav K, Siena S, Takashima A, et al. Trastuzumab deruxtecan in patients with HER2-positive advanced colorectal cancer (DESTINY-CRCO2): primary results from a multicentre, randomised, phase 2 trial. *Lancet Oncol* 2024;25:1147. [PubMed](#) Metastatic disease
57. **CodeBreak 300 OS:** Pietrantonio F, Salvatore L, Esaki T, et al. Overall survival analysis of the phase III CodeBreak 300 study of sotorasib plus panitumumab versus investigator's choice in chemorefractory KRAS G12C colorectal cancer. *J Clin Oncol* 2025;43:2147. [PubMed](#) Metastatic disease
58. **KRYSTAL-1 (adagrasib):** Yaeger R, Weiss J, Pelster MS, et al. Adagrasib with or without cetuximab in colorectal cancer with mutated KRAS G12C. *N Engl J Med* 2023;388:44. [PubMed](#) Metastatic disease
59. **Dostarlimab dMMR rectal (Cercek):** Cercek A, Lumish M, Sinopoli J, et al. PD-1 blockade in mismatch repair-deficient, locally advanced rectal cancer. *N Engl J Med* 2022;386:2363. [PubMed](#) Rectal cancer
60. **RAPIDO:** Bahadoer RR, Dijkstra EA, van Etten B, et al. Short-course radiotherapy followed by chemotherapy before total mesorectal excision (TME) versus preoperative chemoradiotherapy, TME, and optional adjuvant chemotherapy in locally advanced rectal cancer (RAPIDO). *Lancet Oncol* 2021;22:29. [PubMed](#) Rectal cancer
61. **OPRA:** Garcia-Aguilar J, Patil S, Gollub MJ, et al. Organ preservation in patients with rectal adenocarcinoma treated with total neoadjuvant therapy. *J Clin Oncol* 2022;40:2546. [PubMed](#) Rectal cancer
62. **PRODIGE 23:** Conroy T, Bosset JF, Etienne PL, et al. Neoadjuvant chemotherapy with FOLFIRINOX and preoperative chemoradiotherapy for patients with locally advanced rectal cancer (UNICANCER-PRODIGE 23). *Lancet Oncol* 2021;22:702. [PubMed](#) Rectal cancer
63. **PROSPECT:** Schrag D, Shi Q, Weiser MR, et al. Preoperative treatment of locally advanced rectal cancer. *N Engl J Med* 2023;389:322. [PubMed](#) Rectal cancer
64. **Dutch TME:** Kapiteijn E, Marijnen CA, Nagtegaal ID, et al. Preoperative radiotherapy combined with total mesorectal excision for resectable rectal cancer. *N Engl J Med* 2001;345:638. [PubMed](#) Rectal cancer
65. **Stockholm III:** Erlandsson J, Holm T, Pettersson D, et al. Optimal fractionation of preoperative radiotherapy and timing to surgery for rectal cancer (Stockholm III). *Lancet Oncol* 2017;18:336. [PubMed](#) Rectal cancer
66. **German CAO/ARO/AIO-94:** Sauer R, Becker H, Hohenberger W, et al. Preoperative versus postoperative chemoradiotherapy for rectal cancer. *N Engl J Med* 2004;351:1731. [PubMed](#) Rectal cancer
67. **FACS:** Primrose JN, Perera R, Gray A, et al. Effect of 3 to 5 years of scheduled CEA and CT follow-up to detect recurrence of colorectal cancer: the FACS randomized clinical trial. *JAMA* 2014;311:263. [PubMed](#) Follow-up
68. **COLOFOL:** Wille-Jørgensen P, Syk I, Smedh K, et al. Effect of more vs less frequent follow-up testing on overall and colorectal cancer-specific mortality in patients with stage II or III colorectal cancer: the COLOFOL randomized clinical trial. *JAMA* 2018;319:2095. [PubMed](#) Follow-up
69. **CHALLENGE:** Courneya KS, Vardy JL, O'Callaghan CJ, et al. Structured exercise after adjuvant chemotherapy for colon cancer. *N Engl J Med* 2025;393:13. [PubMed](#) Adjuvant and neoadjuvant therapy for colon cancer
70. **Lynch metachronous risk after surgery (Parry 2011):** Parry S, Win AK, Parry B, et al. Metachronous colorectal cancer risk for mismatch repair gene mutation carriers: the advantage of more extensive colon surgery. *Gut* 2011;60:950. [PubMed](#) Screening and prevention

Educational summary for health professionals, written in our own words from current guidelines and primary trials. No drug doses are given. Verify current guidelines, Indian regulatory approvals and individual patient factors before treatment decisions.